

Flora in ontlasting van gezonde kinderen geboren op verschillende wijzen: Blijvende veranderingen van de darmflora na Sectio Caesarea (keizersnede)

Samenvatting

Achtergrond: Pasgeboren kinderen, geboren in moderne ziekenhuizen, zijn afhankelijk van diverse factoren welke van invloed zijn bij de opbouw van een gezonde darmflora- bijvoorbeeld keizersnede en anti-biotica. Om te onderzoeken wat de tijdsduur van deze externe factoren zijn (langdurig of kortdurig), twee groepen kinderen uit twee verschillende verloskundige technieken, werden onderzocht.

Methode: De bacterieflora in de ontlasting van 64 gezonde kinderen werd onderzocht . 34 kinderen waren ter wereld gekomen via de natuurlijke (vaginale) weg, en 30 d.m.v. een keizersnede waarbij, uit voorzorg, antibiotica aan de moeder werd toegediend voor de ingreep. De flora werd opgekweekt in specifieke en niet specifieke media van kinderen van 3-5, 10, 30, 60 en 180 dagen oud. De gastro-intestinale toestand van de moeder werd twee maanden geobserveerd.

Resultaten: De kolonisatie van darmflora bij kinderen geboren door middel van een keizersnede was vertraagd. De aantallen bifidobacterium-achtige bacteriën en de kolonisatie van Lactobacillus-achtige bacteriën kwamen pas op het niveau van vaginaal¹ geboren kinderen na 1 maand en 10 dagen. Kinderen, geboren via een keizersnede, waren significant minder vaak gekoloniseerd met bacteriën uit de bacteria fragilis groep dan vaginaal geboren kinderen. Op een leeftijd van 6 maanden waren de waarden 36% bij de "keizersnee-kinderen" en 76% bij de kinderen geboren op natuurlijke wijze. Het voorkomen van maagdarmklachten was bij beide groepen gelijk.

Conclusie: Deze studie laat voor het eerst zien dat de ontwikkeling van de darmflora bij kinderen geboren d.m.v. een keizersnede tot 6 maanden verstoord is.

Opmerking: Uit meer onderzoek is gebleken dat het toedienen van specifieke preparaten (pro biotica) de ontwikkeling van de darmflora verbeterd. Daarnaast zijn er aanwijzingen dat de ontwikkeling van de darmflora tot het 2^e levensjaar bepalend is voor de rest van het leven. Een goed ontwikkelde darmflora in kindertijd schijnt minder allergieën en andere klachten te geven.

Een speciaal product (orthiflor start) om de darmflora op te bouwen is te koop in de winkel van deze website (www.homeopathieroermond.nl zie online winkel o.a. aanbiedingen)

¹ = via de natuurlijke weg geboren kinderen

Fecal Microflora in Healthy Infants Born by Different Methods of Delivery: Permanent Changes in Intestinal Flora After Cesarean Delivery

*†Minna-Maija Grönlund, †Olli-Pekka Lehtonen, ‡Erkki Eerola, and *Pentti Kero

*Department of Pediatrics, Turku University Central Hospital, Turku, Finland; †Clinical Microbiology, Turku University Central Hospital, ‡Department of Medical Microbiology, University of Turku, Finland

ABSTRACT

Background: Newborn infants in modern maternity hospitals are subject to numerous factors that affect normal intestinal colonization—for example, cesarean delivery and antimicrobial agents. To study the duration of the effect of external factors on intestinal colonization, two groups of infants with different delivery methods were investigated.

Methods: The fecal flora of 64 healthy infants was studied prospectively. Thirty-four infants were delivered vaginally, and 30 by cesarean birth with antibiotic prophylaxis administered to their mothers before the delivery. The fecal flora was cultured on nonselective and selective media in infants 3 to 5, 10, 30, 60, and 180 days of age. Gastrointestinal signs were recorded daily by the mothers for 2 months.

Results: The fecal colonization of infants born by cesarean delivery was delayed. *Bifidobacterium*-like bacteria and *Lac-*

tobacillus-like bacteria colonization rates reached the rates of vaginally delivered infants at 1 month and 10 days, respectively. Infants born by cesarean delivery were significantly less often colonized with bacteria of the *Bacteroides fragilis* group than were vaginally delivered infants: At 6 months the rates were 36% and 76%, respectively ($p = 0.009$). The occurrence of gastrointestinal signs did not differ between the study groups.

Conclusions: This study shows for the first time that the primary gut flora in infants born by cesarean delivery may be disturbed for up to 6 months after the birth. The clinical relevance of these changes is unknown, and even longer follow-up is needed to establish how long-lasting these alterations of the primary gut flora can be. *JPGN* 28:19-25, 1999. **Key Words:** Bacteroides—Cesarean delivery—Colic—Colonization—Gut—Intestine.

The neonatal period is crucial for intestinal colonization. Born sterile and undergoing colonization within a few days, infants are an open field for colonization by different types of bacteria. Gestational age, type of delivery, and feeding affect the stool flora of young infants (1), but there is little information on which long-term factors influence the bacterial selection process in the gut of young infants (2).

Infants born vaginally apparently acquire their gut flora from maternal vaginal and fecal flora (3), but the environment also contributes. Within maternity wards, nosocomial spread is shown to exist among healthy newborn infants (4,5). For the colonization of infants born by cesarean delivery (CD), the environment is extremely important (6,7). Likewise, if infants are separated from

their mothers for long periods after birth, the environment becomes an important source of colonizing bacteria (5).

Gut colonization is delayed in infants born by CD, and intestinal colonization is consequently abnormal for several weeks (1,8). Most of the studies on gut colonization of infants born by CD have extended only to the first month of life (1,8-10). Overall, however, the long-term stability of the gut flora of newborn infants has been studied systematically in only a few studies (2,11). Such information would clearly be valuable, because changes in the primary gut flora have been associated with gastrointestinal disorders and infantile colic (12-15). We studied the stability of the changes in the fecal flora of newborn infants and the association between certain bacteria and gastrointestinal signs. To study two distinctly different groups of neonates with different gut flora, two study groups were formed: vaginally delivered (VD) infants ($n = 34$) and infants born by CD to mothers who had received antimicrobial prophylaxis ($n = 30$). The fecal flora was recorded for 6 months, and gastrointestinal signs were registered daily for 2 months.

Received December 18, 1997; Revised April 7 and June 26, 1998; accepted July 1, 1998.

Address correspondence and reprint requests to Minna-Maija Grönlund, MD, Department of Paediatrics, Turku University Central Hospital, Kiinamyllynkatu 4-8, FI-20520 Turku, Finland.

MATERIALS AND METHODS

Infants

The study included 64 healthy newborn infants of healthy mothers who delivered at the Department of Obstetrics and Gynecology, University Central Hospital, Turku, Finland. Thirty-four VD infants and 30 infants delivered by elective CD were enrolled between February 1995 and June 1996 after written informed consent had been obtained from their parents. The enrollment rate was two to four children per week, because sampling necessitated that the birth took place on Monday through Thursday. The mothers who delivered by cesarean birth received prophylactically 2 g intravenous ampicillin 2 hours before the operation. None of the mothers had received any antimicrobial agents within the month that preceded the delivery. After delivery, the newborns were admitted randomly to one of the two maternity wards for healthy newborn infants.

Fecal Samples and Bacterial Cultures

Fecal samples were taken when the infants were 3 to 5 days old (taken at the hospital) and 10, 30, 60, and 180 days old (taken at home). The specimens were collected in plastic containers. If not cultured immediately, the samples were stored at 4°C. The specimens obtained at home were taken to the laboratory by the parents. The mean storage time of the specimens was 10.7 hours (range, 0.5–34.5 hours).

An approximate 300-mg portion of the specimen was weighed, diluted, and homogenized in fastidious anaerobe broth (Lab M, Bury, UK). Serial 100-fold dilutions were made in the same broth. Duplicate samples of 10 µl of each dilution were cultured on a variety of nonselective and selective media (Table 1).

The MacConkey plates were incubated at 35°C in ambient air for 24 hours. All other cultures were incubated in anaerobic jars with gas-generating kits (Anero Gen, AN35; Oxoid, Basingstoke, UK) at 35°C for 48 hours (fastidious anaerobe agar [FAA], *Bacteroides* bile esculin agar [BBE] and *Clostridium perfringens* selective agar), or for 72 hours (Rogosa and modified Petuely's agars). The number of colonies from two parallel plates

was counted from a dilution yielding 30 to 300 colony-forming units (CFU)/plate, and the average was recorded.

The total number of colonies was counted and recorded from MacConkey and FAA agars. The number of all different types of colonies was counted from BBE, Rogosa, and modified Petuely's agars and the number of distinctive black colonies from *Clostridium perfringens* agar. All different types of colonies from BBE, Rogosa, and modified Petuely's agars and distinctive black colonies from *Clostridium perfringens* agar were subcultured for aerotolerance testing and were Gram stained.

Anaerobic gram-negative, esculin hydrolysing rods growing on BBE agar were recorded as bacteria of the *Bacteroides fragilis* group. Anaerobic and aerotolerant nonbranching, gram-positive rods with parallel sides from Rogosa agar were recorded as *Lactobacillus*-like bacteria (LLB). It has been shown that all colonies growing well on Rogosa agar may be considered lactic acid bacteria; some enterococci and pediococci may show reduced growth (16). Anaerobic and aerotolerant gram-positive rods from modified Petuely's agars were recorded as *Bifidobacterium*-like bacteria (BLB). Modified Petuely's agar is highly selective and efficient for detection of *Bifidobacterium* from fecal samples. In a previous work by Tanaka and Mutai (17) 175 fecal strains were detected from the modified Petuely's medium; from these 94% were bifidobacteria, 3% eubacteria, and 3% peptostreptococci.

Clostridium perfringens was identified by its distinct colony and Gram-stain morphology, by its anaerobic nature in aerotolerance testing, and by positive results in the reversed CAMP test (18). The bacterial counts were expressed as the log₁₀ of colony-forming units per gram of wet weight of feces.

Clinical Symptoms

The mothers kept a diary for 2 months of their infants' daily bowel habits, abdominal distension, flatulence, normal and colicky crying, use of antimicrobial agents, and feedings. Crying was defined as colicky if it was a distinctive pain cry, and the infant was difficult to console (19). Abdominal distension and flatulence were given a daily score by the following scale: no signs, a few signs, moderate signs, or heavy signs.

TABLE 1. Culture media and dilutions

| Medium | Bacteria | Dilutions |
|--|--|--|
| MacConkey agar (CM7, Oxoid, Basingstoke, UK) | Aerobic enteric bacteria ^a | 3-day, 10-day, 30-day samples ^b 10 ⁵ –10 ⁹ 60-day, 180-day samples 10 ⁷ –10 ¹¹ |
| Fastidious anaerobe agar (Lab M, Bury, UK) | Total count ^c | 3-day, 10-day, 30-day samples 10 ⁵ –10 ⁹ 60-day, 180-day samples 10 ⁷ –10 ¹¹ |
| <i>Bacteroides</i> bile esculin agar (Reference 21) | <i>Bacteroides fragilis</i> group | 3-day, 10-day, 30-day samples 10 ³ –10 ⁷ 60-day, 180-day samples 10 ⁵ –10 ⁹ |
| <i>Clostridium perfringens</i> selective agar (CM 543, Oxoid) | <i>Clostridium perfringens</i> | 3-day, 10-day, 30-day samples 10 ³ –10 ⁷ 60-day, 180-day samples 10 ⁵ –10 ⁹ |
| Rogosa agar (CM 627, Oxoid) | <i>Lactobacillus</i> -like bacteria ^d | 3-day sample 10 ³ –10 ⁷ 10-day, 30-day, 60-day, 180-day samples 10 ⁵ –10 ⁹ |
| Modified Petuely's agar (Reference 17) | <i>Bifidobacterium</i> -like bacteria | 3-day sample 10 ³ –10 ⁷ 10-day, 30-day, 60-day, 180-day samples 10 ⁵ –10 ⁹ |

^a Includes enterobacteria, staphylococci, enterococci.

^b 3–5-day, 10-day, and 30-day samples.

^c Includes facultative and anaerobic bacteria.

^d Includes gram-positive, non-branching rods with parallel sides.

Statistical Analysis

The results were analysed statistically by Fisher's exact test (to compare the number of infants colonized at each time point), Mann-Whitney's rank sum test (to compare bacteria counts of colonized infants at 3 days and 6 months of age), and analysis of variance of repeated measurements (to compare the gastrointestinal signs). A commercial software program (Statistica, version 5.0; Stat Soft, Tulsa, OK, U.S.A.) was used for these calculations. $P < 0.05$ was considered statistically significant.

ETHICAL CONSIDERATIONS

The study was approved by the joint Committee of Ethics of the Turku University and Turku University Central Hospital.

RESULTS

The study groups differed slightly by gestational age (VD group, 40 weeks; CD group, 39 weeks; $p = 0.04$) but the birth weights did not differ statistically (VD group, 3577 g; CD group, 3572 g; $p = 0.96$). Special attention was paid to the method of feeding the infants. The use of formula was recorded carefully by the mothers. The proportion of infants exclusively breast-fed at 2 months of age or partly breast-fed at 6 months of age did not differ between the study groups ($p = 0.2$ and 0.8 , respectively). Eleven infants received antimicrobial agents during follow-up. All of these therapies were administered when the infants were older than 2 months, and in these infants the subsequent 6-month fecal samples were excluded from the analysis. (Table 2)

TABLE 2. Study subjects

| | Vaginally delivered (n = 34) | Cesarean delivery (n = 30) | p |
|---|---------------------------------|-------------------------------|-------------------|
| Mean birth weight in grams (range) | 3577 (2850-4480) | 3572 (2640-4320) | 0.96 ^a |
| Mean gestational age in weeks (range) | 40 (37-42) | 39 (37-42) | 0.04 ^a |
| Number of boys | 16 | 12 | 0.6 ^b |
| Number of infants exclusively breast-fed for 2 months | 19 | 22 | 0.2 ^b |
| Number of infants partly breast-fed for 6 months ^c | 17 | 13 | 0.8 ^b |
| Number of infants receiving antibiotic regimen during follow-up | 3 | 8 | 0.1 ^b |

^a Two-sample t-test.

^b Fisher's exact test.

^c Information missing from two vaginally delivered infants and from three cesarean births.




FIG. 1. The percentage of *Bifidobacterium*-like bacteria (BLB) colonization in infants aged 3, 10, 30, 60, and 180 days born vaginally and by cesarean delivery. * $p < 0.001$.

Colonization Rates

With the exception of one infant (in the CD group) who was culture-negative at 3 days of age, all infants were colonized with aerobic enteric bacteria in every culture. The colonization rates of BLB and LLB were lower in the CD group than in the VD group after birth. The colonization rate of BLB coincided in these groups by 1 month and of LLB by 10 days of age (Figs. 1 and 2). The colonization rate of LLB in the CD group even exceeded that of the VD group in infants 2 and 6 months of age (Fig. 2). The *Clostridium perfringens* colonization rate was statistically higher in the CD group than in the VD group at 1 month of age (57% vs. 17%; $p = 0.003$; Fig. 3).

The colonization rates of bacteria within the *Bacteroides fragilis* group differed most markedly of all between the study groups. The *Bacteroides* colonization rate ranged from 52% to 79% in the VD group (Fig. 4). Only one of the infants in the CD group was *Bacteroides*-positive at 3 days of age. After that, *Bacteroides* was not recovered in any of the samples from the infants in the CD group before the age of 2 months. In infants 6 months of age, the colonization rate was still statistically lower in the CD group than in the VD group (36% vs. 76%; $p = 0.009$; Fig. 4).

Colonization Levels

The colonization levels of the different bacteria of the colonized infants are shown in Table 3. The total bacterial counts were significantly lower in the CD group in infants 3 days of age ($p = 0.005$) and in those 6 months of age ($p = 0.03$). The aerobic enteric bacterial counts did not differ between the groups. Infants in the CD group had a significantly lower level of BLB when they were 3 days old ($p = 0.005$) but not when they were 6 months old ($p = 0.5$). There were no differences in the




FIG. 2. The percentage of *Lactobacillus*-like bacteria (LLB) colonization in infants aged 3, 10, 30, 60, and 180 days born vaginally and by cesarean delivery.

LLB counts or in the *Clostridium perfringens* counts between the groups. The amounts of *Bacteroides fragilis* bacteria did not differ between the groups of infants with colonization at 6 months ($p = 0.3$).

Gastrointestinal Signs

Two mothers in both groups did not complete the follow-up sheets. No statistically significant differences were found between the study groups in the scores of abdominal distension, flatulence, or in the amount of colicky crying. Infantile colic, according to the definition of Wessel et al. (20), was detected in three of the VD infants and in none of the CD infants ($p = 0.2$; Fisher's exact test).

DISCUSSION

This study shows that the fecal flora of infants born by CD with prophylactic antibiotics administered to the




FIG. 3. The percentage of *Clostridium perfringens* colonization in infants aged 3, 10, 30, 60, and 180 days born vaginally and by cesarean delivery. * $p = 0.003$.




FIG. 4. The percentage of *Bacteroides fragilis* group colonization in infants aged 3, 10, 30, 60, and 180 days born vaginally and by cesarean delivery. * $p < 0.001$, † $p = 0.009$.

mother is very different from that of infants delivered vaginally. The greatest differences were seen in the bacteria of the *Bacteroides fragilis* group, which is in agreement with studies in infants born by CD; in earlier studies the duration of follow-up, however, was only 10 to 60 days (1,8,10). In the present study, no permanent colonization with *Bacteroides fragilis* group bacteria was found in the CD group before the infants were 2 months of age. Still, in infants 6 months of age in the CD group, the colonization rate of the *Bacteroides fragilis* group was only half that of infants in the VD group (36% and 76%, respectively; $p = 0.009$). Similarly, Bennet and Nord (8) were unable to detect *Bacteroides* bacteria 3 to 8 weeks after birth in term and preterm infants born by CD.

We used selective culture media to describe intestinal flora in this study. The counts detected from these selective culture media represent mainly the bacterial genera level but cannot detect the absolute counts of different bacterial genera or species, except those of *Clostridium perfringens*, which were further identified. Nevertheless, this method can identify differences between the delivery groups and the changes taking place in intestinal flora after birth, which was the main purpose of the current study. Furthermore, the main differences in the fecal flora between the delivery groups were found in *Bacteroides fragilis* group bacteria that were detected from the BBE media, which is highly selective for the *Bacteroides fragilis* group (21).

The long-lasting changes seen in the primary gut flora of infants born by CD could be the result of one or both of the two abnormal components of their birth: CD itself or the prophylactic ampicillin administered to the mother 2 hours before the elective CD. Ampicillin is very poorly protein-bound and crosses the placenta readily; maternal and fetal serum ampicillin levels equilibrate within 1 hour after intravenous administration (22). The decline in BLB could be explained by the intravenous ampicillin that the infants were exposed to before delivery, but fecal

TABLE 3. Median and range of fecal bacterial counts of colonized infants at 3, 10, 30, 60, and 180 days of age

| Age (days) | Vaginal delivery | | | | | Cesarean delivery | | | | |
|-----------------------------------|-------------------|----------|-----------|-----------|-------------------|-------------------|----------|-----------|-----------|-----------|
| | 3 | 10 | 30 | 60 | 180 | 3 | 10 | 30 | 60 | 180 |
| Bacteria | | | | | | | | | | |
| Total | 10.6 ^a | 11.0 | 11.1 | 11.3 | 11.4 ^c | 10.1 | 11.1 | 11.1 | 11.2 | 11.1 |
| range | 10.1-11.8 | 9.3-12.1 | 10.2-11.9 | 10.1-13.5 | 9.9-13.5 | 7.0-11.9 | 9.5-11.8 | 10.3-11.7 | 10.4-13.7 | 10.1-13.1 |
| n | 34 | 31 | 33 | 33 | 29 | 30 | 28 | 29 | 29 | 22 |
| Aerobic enteric | 9.6 | 9.6 | 9.6 | 9.5 | 9.4 | 9.8 | 9.9 | 9.9 | 9.8 | 9.6 |
| range | 6.7-10.6 | 8.1-10.8 | 7.3-10.6 | 6.6-12.4 | 7.5-11.7 | 6.6-10.9 | 8.8-12.0 | 7.2-11.0 | 8.5-13.1 | 8.1-11.6 |
| n | 34 | 30 | 33 | 33 | 29 | 29 | 27 | 29 | 29 | 22 |
| <i>Bacteroides fragilis</i> group | 8.7 | 9.3 | 9.4 | 9.5 | 9.6 | 6.9 | — | — | 8.1 | 8.7 |
| range | 4.8-10.2 | 4.0-10.4 | 5.5-10.6 | 5.5-11.1 | 5.5-12.3 | — | — | — | 5.5-10.5 | 6.8-11.2 |
| n | 22 | 16 | 33 | 26 | 22 | 1 | 0 | 0 | 4 | 8 |
| <i>Lactobacillus</i> -like | 7.8 | 7.7 | 8.3 | 9.3 | 9.6 | 6.6 | 9.4 | 9.4 | 8.8 | 8.9 |
| range | 6.4-10.2 | 6.8-10.4 | 6.6-9.8 | 7.4-10.9 | 5.8-10.1 | — | 6.5-10.5 | 7.4-10.3 | 5.7-10.3 | 6.8-10.1 |
| n | 6 | 7 | 10 | 11 | 10 | 1 | 7 | 11 | 18 | 13 |
| <i>Bifidobacterium</i> -like | 10.2 ^b | 10.7 | 10.8 | 10.9 | 10.9 | 8.2 | 10.7 | 10.8 | 10.9 | 10.8 |
| range | 7.8-11.1 | 7.5-11.4 | 6.4-11.6 | 8.9-11.8 | 9.0-12.0 | 5.1-10.8 | 5.1-11.8 | 7.5-11.5 | 7.2-11.6 | 8.9-12.1 |
| n | 28 | 30 | 30 | 31 | 27 | 11 | 15 | 24 | 24 | 20 |
| <i>Clostridium perfringens</i> | 6.7 | 6.0 | 7.4 | 8.2 | 6.6 | 7.5 | 6.9 | 8.2 | 6.9 | 7.0 |
| range | 3.8-8.7 | 4.4-8.0 | 6.8-9.0 | 5.8-9.0 | 5.8-8.5 | 7.52-7.53 | 4.1-9.1 | 4.5-9.5 | 5.8-10.0 | 5.5-9.0 |
| n | 5 | 4 | 5 | 5 | 7 | 2 | 7 | 16 | 9 | 4 |

Counts expressed as log₁₀ colony-forming units per gram wet weight of feces.

n, number of colonized infants. P are vaginal delivery versus cesarean delivery.

^a p = 0.005.

^b p = 0.005.

^c p = 0.03.

LBL should not decline with ampicillin therapy, at least this is not the case in older children (10 months to 12 years). Also, only a minor decline is detected in fecal *Bacteroides* sp. counts in these older children and the *Bacteroides* sp. counts return to normal stage within only 3 to 6 days after cessation of the ampicillin therapy (23). Thus, it seems unlikely that the long-term changes recorded in the fecal flora of the infants in the CD group in our study could be explained by the administration of ampicillin before delivery. Rather, previous reports imply that this phenomenon is caused by CD itself (1,8). Still, ampicillin could have more profound effects when administered immediately after the birth because the gut is empty, and the drug can select the primary colonizing bacteria.

The changes that took place in the fecal bacterial flora could not be associated with gastrointestinal signs. Any signs were meticulously recorded daily by the mothers for 2 months. In a previous study, *Clostridium perfringens* has been associated with an increased incidence of gastrointestinal signs, such as flatulence, distended abdomen, foul-smelling stools, diarrhea, and blood in stools (12). In the present study, the infants born by CD had a higher colonization rate of *Clostridium perfringens* than the VD group of infants at 1 month (57% and 17%, respectively). Even at this time point, the scores of gastrointestinal signs did not differ between the study groups. The inconsistency of the results between these two studies might be explained by the differences between the study populations. In the previous study most of the infants were treated in the intensive care unit and were

preterm infants, whereas in the present study all infants were healthy full-term neonates.

There is great variation among the reports of the predominant bacteria in the fecal flora of breast-fed newborns. Some researchers have found bifidobacteria to predominate and *Bacteroides* to be scarce (2,24,25), whereas others have found the opposite (26,27). However, the predominant bacterium has been classified differently among the studies. Some investigators define the predominant bacterium as the most frequently occurring bacteria (26) and some as the bacterium with the highest counts in fecal samples (28). In the present study, 19 of the 34 VD infants were exclusively breast-fed for 2 months. Among all the VD children, the most frequent bacteria during the first 2 months of life were BLB (85-97%), although *Bacteroides fragilis* group bacteria were also common (52-79%). The highest colony counts were encountered for BLB (10.2-10.9 log₁₀ CFU/g wet weight of feces). The *Bacteroides* colony counts were lower (8.7-9.5 log₁₀ CFU/g). These results agree with those in a recent study from Germany (28).

The present study is the first one to show that the changes in the primary intestinal flora of infants born by CD to mothers who have received antimicrobial prophylaxis last for no less than 6 months after birth, maybe longer. We do not know how long these abnormalities in the fecal flora last on the whole, but the primary quality and quantity of colonization of the gut seems to be critical in the selection process between different genera of bacteria. Some bacteria such as *Bacteroides* are not as easily accessible to the gut of infants delivered the under

sterile conditions of CD. Other, perhaps more aerotolerant, bacteria may take over in the intestine and inhibit the subsequent colonization of the gut by *Bacteroides*. This phenomenon is the generally known as interbacterial inhibition (29). Apparently, the bacterial predominances are not readily subject to change under normal domestic conditions after the first months of the colonization process.

Normal intestinal flora has immunostimulatory functions, as has been demonstrated in numerous animal studies (30-32). Mucosal IgA plasma cells are especially scarce in germ-free animals (33). Additionally, when probiotic bacteria belonging to the normal intestinal flora have been administered orally to children in association with diarrhea or mucosal vaccination, an increase in antigen-specific and nonspecific IgA and IgM responses has been detected (34,35). Secretory IgA and IgM are the main humoral mediators of mucosal immunity in cooperation with a variety of innate protective mechanisms. Well-functioning mucosal immunity is a prerequisite for health, because the mucosal surfaces are favored as portals of entry by most infectious agents, allergens, and carcinogens (36). Further research is under way to determine whether the delay found in intestinal colonization in these infants born by CD has any effect on the development of the gut-associated immune system.

Acknowledgement: The authors thank Mr. Hans Helenius, M.Sc., Department of Biostatistics, University of Turku, for his help with the statistical analysis; and the nurses of the Department of Obstetrics and Gynecology, and Mrs. Satu Ekblad, Department of Pediatrics, for their invaluable help in performing this study.

Supported in part by The South-West Finnish Fund of Neonatal Research.

REFERENCES

- Long SS, Swenson RM. Development of anaerobic fecal flora in healthy newborn infants. *J Pediatr* 1977;91:298-301.
- Stark PL, Lee A. The microbial ecology of the large bowel of breast-fed and formula-fed infants during the first year of life. *J Med Microbiol* 1982;15:189-203.
- Tannock GW, Fuller R, Smith SL, Hall MA. Plasmid profiling of members of the family Enterobacteriaceae, lactobacilli, and bifidobacteria to study the transmission of bacteria from mother to infant. *J Clin Microbiol* 1990;28:1225-8.
- Delmeé M, Verellen G, Avesani V, Francois G. Clostridium difficile in neonates: Serogrouping and epidemiology. *Eur J Pediatr* 1988;147:36-40.
- Murono K, Fujita K, Yoshikawa M, et al. Acquisition of nonmaternal Enterobacteriaceae by infants delivered in hospitals. *J Pediatr* 1993;122:120-5.
- Fryklund B, Tullus K, Berglund B, Burman LG. Importance of the environment and the faecal flora of infants, nursing staff and parents as sources of gram-negative bacteria colonizing newborns in three neonatal wards. *Infection* 1992;20:253-7.
- Lennox-King SM, O'Farrell SM, Bettelheim KA, Shooter RA. Escherichia coli isolated from babies delivered by caesarean section and their environment. *Infection* 1976;4:139-45.
- Bennet R, Nord CE. Development of the faecal anaerobic microflora after caesarean section and treatment with antibiotics in newborn infants. *Infection* 1987;15:332-6.
- Hall MA, Cole CB, Smith SL, Fuller R, Rolles CJ. Factors influencing the presence of faecal lactobacilli in early infancy. *Arch Dis Child* 1990;65:185-8.
- Neut C, Bezirtzoglou E, Romond C, Beerens H, Delcroix M, Noel AM. Bacterial colonization of the large intestine in newborns delivered by caesarean section. *Zentralbl Bakteriell Mikrobiol Hyg A* 1987;266:330-7.
- Ellis-Pegler RB, Crabtree C, Lambert HP. The faecal flora of children in the United Kingdom. *J Hyg Lond* 1975;75:135-42.
- Ahtonen P, Lehtonen OP, Kero P, Eerola E, Hartiala K. Clostridium perfringens in stool, intrapartum antibiotics and gastrointestinal signs in a neonatal intensive care unit. *Acta Paediatr* 1994; 83:389-90.
- Lehtonen L, Korvenranta H, Eerola E. Intestinal microflora in colicky and noncolicky infants: Bacterial cultures and gas-liquid chromatography. *J Pediatr Gastroenterol Nutr* 1994;19:310-4.
- Miller JJ, McVeagh P, Fleet GH, Petocz P, Brand JC. Breath hydrogen excretion in infants with colic. *Arch Dis Child* 1989;64: 725-9.
- Moore DJ, Robb TA, Davidson GP. Breath hydrogen response to milk containing lactose in colicky and noncolicky infants. *J Pediatr* 1988;113:979-84.
- The Oxoid Manual*. 7th ed. Basingstoke, UK: Unipath Ltd., 1995: 2-186.
- Tanaka R, Mutai M. Improved medium for selective isolation and enumeration of Bifidobacterium. *Appl Environ Microbiol* 1980;40: 866-9.
- Buchanan AG. Clinical laboratory evaluation of a reverse CAMP test for presumptive identification of Clostridium perfringens. *J Clin Microbiol* 1982;16:761-2.
- Lester BM, Boukydis CFZ, Garcia-Coll CT, Hole WT. Colic for developmentalists. *Infant Mental Hlth J* 1990;11:321-33.
- Wessel MA, Cobb JC, Jackson EB, Harris GSJ, Detweiler AC. Paroxysmal fussing in infancy, sometimes called "colic." *J Pediatr* 1954;14:421-34.
- Livingston SJ, Kominos SD, Yee RB. New medium for selection and presumptive identification of the Bacteroides fragilis group. *J Clin Microbiol* 1978;7:448-53.
- Bray RE, Boe RW, Johnson WL. Transfer of ampicillin into fetus and amniotic fluid from maternal plasma in late pregnancy. *Am J Obstet Gynecol* 1966;96:938-42.
- Sakata H, Fujita K, Yoshioka H. The effect of antimicrobial agents on fecal flora of children. *Antimicrob Agents Chemother* 1986;29: 225-9.
- Roberts AK, Chierici R, Sawatzki G, Hill MJ, Volpato S, Vigi V. Supplementation of an adapted formula with bovine lactoferrin: 1. Effect on the infant faecal flora. *Acta Paediatr* 1992;81:119-24.
- Yoshioka H, Iseki K, Fujita K. Development and differences of intestinal flora in the neonatal period in breast-fed and bottle-fed infants. *Pediatrics* 1983;72:317-21.
- Lundequist B, Nord CE, Winberg J. The composition of the faecal microflora in breastfed and bottle fed infants from birth to eight weeks. *Acta Paediatr Scand* 1985;74:45-51.
- Simhon A, Douglas JR, Drasar BS, Soothill JF. Effect of feeding on infants' faecal flora. *Arch Dis Child* 1982;57:54-8.
- Kleessen B, Bunke H, Tovar K, Noack J, Sawatzki G. Influence of two infant formulas and human milk on the development of the faecal flora in newborn infants. *Acta Paediatr* 1995;84:1347-56.
- Sprunt K, Redman W. Evidence suggesting importance of role of interbacterial inhibition in maintaining balance of normal flora. *Ann Intern Med* 1968;68:579-90.
- Crabbé PA, Bazin H, Eysen H, Heremans JF. The normal microbial flora as a major stimulus for proliferation of plasma cells synthesizing IgA in the gut. The germ-free intestinal tract. *Int Arch Allergy Appl Immunol* 1968;34:362-75.
- Moreau MC, Ducluzeau R, Guy-Grand D, Muller MC. Increase in

- the population of duodenal immunoglobulin A plasmocytes in axenic mice associated with different living or dead bacterial strains of intestinal origin. *Infect Immun* 1978;21:532-9.
32. Shroff KE, Meslin K, Cebra JJ. Commensal enteric bacteria engender a self-limiting humoral mucosal immune response while permanently colonizing the gut. *Infect Immun* 1995;63:3904-13.
 33. Crabbé PA, Nash DR, Bazin H, Eyssen H, Heremans JF. Immunohistochemical observations on lymphoid tissues from conventional and germ-free mice. *Lab Invest* 1970;22:448-57.
 34. Kaila M, Isolauri E, Soppi E, Virtanen E, Laine S, Arvilommi H. Enhancement of the circulating antibody secreting cell response in human diarrhea by a human Lactobacillus strain. *Pediatr Res* 1992; 32:141-4.
 35. Isolauri E, Joensuu J, Suomalainen H, Luomala M, Vesikari T. Improved immunogenicity of oral D × RRV reassortant rotavirus vaccine by Lactobacillus casei GG. *Vaccine* 1995;13:310-2.
 36. Brandtzaeg P. Molecular and cellular aspects of the secretory immunoglobulin system. *APMIS* 1995;103:1-19.

In the Next Issue

Original Articles

Sarcosidase Therapy for Congenital Sucrase-Isomaltase Deficiency

William R. Treem, Luanne McAdams, Lesley Stanford, Geraldine Kastoff, Christopher Justinich, and Jeffrey Hyams

Helicobacter pylori Infection in Children with Celiac Disease: Prevalence and Clinicopathologic Features

F. Lizza, M. Mancuso, M. Imeneo, L. Mesuraca, A. Contaldo, L. Giancotti, A. La Vecchia, C. Docimo, L. Pensabene, P. Strisciuglio, F. Pallone, and S. Guandalini

Cisapride in Pediatric Chronic Constipation Management

Issam M. Halabi

Home Parenteral Nutrition in Children: The Polish Experience

Janusz Ksiazek, Malgorzata Lyszkowska, Jaroslaw Kierkus, Krzysztof Bogucki, Anna Ratyńska, Bozena Tondys, and Jerzy Socha

Case Report

Congenital Microvillus Atrophy in a Girl with Autosomal Dominant Hypochondroplasia

Peter Heinz-Erian, Heinrich Schmidt, Martine Le Merrer, Alan D. Phillips, Wieland Kiess, and Hans-Beat Hadorn